1 Rebound effects may undermine benefits of upcycling food waste and

- 2 food processing by-products as animal feed in China
- 3
- 4 Weitong Long^{1,2}, Xueqin Zhu^{1*}, Hans-Peter Weikard¹, Oene Oenema^{2,3}, Yong Hou^{2*}
- 5
- 6 ¹Environmental Economics and Natural Resources Group, Wageningen University, Hollandseweg
- 7 1, 6706 KN Wageningen, The Netherlands
- 8 ²State Key Laboratory of Nutrient Use and Management, College of Resources and Environmental
- 9 Science, China Agricultural University, 100193 Beijing, China
- ³Wageningen Environmental Research, 6708 PB Wageningen, The Netherlands

- * Corresponding author at: Wageningen University, 6706 KN Wageningen, The Netherlands; China
- 13 Agricultural University, 100193, Beijing, China.
- 14 E-mail addresses: xueqin.zhu@wur.nl (X. Zhu); yonghou@cau.edu.cn (Y. Hou).

Abstract

Upcycling food waste and food processing by-products as animal feed could reduce environmental impacts of livestock production, but rebound effects, where lower feed costs lead to livestock production expansion, may diminish these benefits. Using an integrated environmental-economic model, we assessed the global impacts of upcycling food waste and food processing by-products in China's monogastric livestock production. We found that the upcycling increased monogastric livestock production by 23-36% and raised Chinese economy-wide acidification emissions by 2.5-4.0%. Greenhouse gas emissions decreased by 0.5-1.4% through less food waste and food processing by-products in landfills and incinerators and contraction of the non-food sectors. This upcycling, accompanied by resource reallocation across the whole economy, enhanced food security and had significant knock-on effects beyond the agricultural sectors, thereby influencing sectoral employment, gross domestic product, and household welfare. Implementing appropriate emission taxes provides an opportunity to absorb the rebound effects on emissions but may negatively affect food security indicators and shift emission-intensive sectors from China to its trading partners, depending on the height of the taxes. Our study, thus, supports policy design aimed at achieving environmental sustainability and food security.

Keywords

- circular food system; food waste; food security; environmental impacts; environmental-economic
- 34 modelling; rebound effects.

Main

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

Animal-sourced food (ASF), such as meat, milk, and eggs, is the main contributor to the environmental impacts of food systems. The surge in demand for ASF, driven by population growth, and increased prosperity and urbanization, 1,2 is expected to double by 2050, especially in developing countries ³. This surge in livestock production has exacerbated food-feed competition and significantly contributes to the exceedance of the planetary boundaries (PBs) for nitrogen (N), phosphorus (P) and greenhouse gas (GHG) emissions. Currently, 70% of global agricultural land is used for producing animal feed 4, and global livestock production accounts for 13-18% of the total anthropogenic GHG emissions ⁵, 40% of the ammonia (NH₃) and nitrous oxide (N₂O) emissions ⁶, and around 24% of N and 55% of P losses to water bodies 7. It has been argued that the global 1.5°C climate target cannot be achieved without mitigating emissions from food systems ⁸. Global food waste has risen from 1.3 to 1.6–2.5 billion tons in recent years despite efforts to reduce food waste 9. A large proportion of food waste ends up in landfills or incinerators, exacerbating GHG emissions and associated climate change 10. Upcycling food waste and food processing byproducts (also called "low-opportunity-cost feed products (LCFs)"), as animal feed is, thus, crucial for reducing environmental impacts and building more circular food systems 11, as it offers a pathway to mitigate land-related pressures 12, alleviate the food-feed competition 11, and reduce emissions from food systems and improper food waste disposal ¹³. This is because food waste and food processing by-products typically compete less for land and natural resources than humanedible feed crops ¹¹⁻¹³. Increased utilisation of food waste and food processing by-products as feed may also contribute to achieving Sustainable Development Goals (SDGs), including SDG 2 (zero hunger), SDG 6 (clean water and sanitation), SDG 12 (responsible consumption and production), SDG 13 (climate action), and SDG 15 (life on land) ¹⁴. While many studies acknowledge the environmental benefits of upcycling food waste and food processing by-products as animal feed, significant gaps remain in the existing literature, particularly in three critical areas. First, previous studies ¹¹⁻¹³ employing linear optimization models to evaluate the environmental impacts of this circular transition may have overestimated the environmental benefits by disregarding "rebound effects" (also known as "Jevons paradox") 15. The rebound effect,

where lower feed costs lead to livestock production expansion, may diminish the environmental benefits of feeding animals with food waste and food processing by-products. Second, the "rebound effect" phenomenon has been extensively studied in energy systems 16,17, but studies of its implications in food systems are largely lacking. Although previous studies have explored rebound effects related to a global dietary shift towards plant-based food ¹⁸ and halving food loss and waste ¹⁹, there is still limited understanding of the rebound effect of upcycling food waste and food processing by-products as animal feed. Third, strateiges to absorb these negative rebound effects resulting from upcycling food waste and food processing by-products as animal feed have not yet been explored. Implementing emissions taxes is considered as an effective policy instrument to identify the most cost-effective mitigation pathway for achieving a given emission mitigation target ²⁰⁻²². For example, many countries, such as the United States, France, Canada, and New Zealand, have implemented various forms of carbon taxes to mitigate GHG emissions ²³. China has committed to tackling both global environmental challenges, such as reducing GHG emissions through its pledge for carbon neutrality by 2060 under the Paris Agreement ^{24,25}, as well as addressing local environmental pollution, such as nitrogen oxides (NO_x), and sulphur dioxide (SO₂) emissions, to meet the reduction targets set in the "14th Five-Year Plan" 26. The Chinsese government recently released a national plan to reduce concentrate feedstuffs such as soybean and maize in pig and chincken production sectors through improved feeding strategies including the upcycling food waste and food processing by-products as animal feed. Evidently, there is a great need to better understand the potential rebound effects that may influence the expected benefits of upcycling food waste and food processing by-products as animal feed, before this action plan is widely implemented in China. In this study, we tried to fill these gaps and thereby contribute to the existing literature by using an integrated environmental-economic applied general equilibrium (AGE) modelling approach to assess the environmental and economic consequences of upcycling food waste and food processing by-products in China's monogastric livestock production as feed, in a global context. Next, we explored how implementing economy-wide emissions taxes could absorb rebound effects of this upcycling while safeguarding food security. We focused on China for our study because it is the world's largest animal producer, accounting for 46%, 34%, and 13% of global pork, egg, and poultry

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

meat production in 2018, respectively ²⁷. Furthermore, around 27% of of food produced for human consumption is lost or wasted in China ²⁸, implying a great opportunity to upcycle the discared food waste as feed. In addition, the Chinese government has proposed to lower the agricultural product processing loss to below 3% by 2035 ²⁹, and to substitute human-edible feed ingredients, such as soybeans and maize, in animal feed with food processing by-products ³⁰. Thus, we considered food waste (cereal grains waste, vegetables & fruits waste, roots & tubers waste, and oilseeds & pulses waste) and food processing by-products (cereal bran, alcoholic pulp, and oil cakes). We addressed three main research questions. First, how will an increased utilisation of food waste and food processing by-products as feed influence livestock production, food supply, and other sectors in China and its main food and feed trading partners (MTP, including Brazil, the United States, and Canada)? Second, how will an increased utilisation of food waste and food processing by-products influence economy-wide emissions of GHGs, acidification pollutants, and eutrophication pollutants, as well as food security indicators (i.e., average food price, food affordability, population at risk of hunger, and food availability)? Third, how will emission taxes absorb rebound effects of this upcycling while safeguarding food security? We examined five scenarios: (i) the baseline (S0) scenario represents the economies of China and MTP in 2014; (ii) scenario 1 (S1) involves upcycling partial use of food waste and food processing by-products (54% of food waste and 100% of food processing by-products) as feed for monogastric livestock production in China; (iii) scenario 2 (S2) involves upcycling full use of food waste and food processing by-products (100% of food waste and 100% of food processing by-products) as feed for monogastric livestock production in China; (iv) scenario 3 (S3 = S1 + A modest emission)mitigation target) entails implementing economy-wide emission taxes to ensure that emissions of GHGs, acidification pollutants, and eutrophication pollutants in both China and MTP do not exceed their baseline (S0) levels; (v) scenario 4 (S4 = S1 + An ambitious emission mitigation target) entails implementing economy-wide emission taxes to meet China's and MTP's annual GHG mitigation targets under the Intended Nationally Determined Contributions (INDC) of the Paris Agreement ^{24,25}, while also addressing China's emission reduction goals for acidification and eutrophication pollutants in line with the "14th Five-Year Plan" ²⁶. The levels of upcycling partial and full use of food waste and food processing by-products as animal feed is estimated using calculations from

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

Fang, et al. ¹², who determine that the maximum utilisation rate of food waste with high moisture content in China is 54% when cross-provincial transportation of food waste is not allowed. When substituting primary feed (i.e., feed crops and compound feed) in animal diets with food waste and food processing by-products, the total protein and total energy supplies per unit of animal output were kept constant in all scenarios (See Supplementary Table 1).

China produced about 104 Tg of monogastric livestock products (pork: 57 Tg; poultry meat: 18 Tg;

Results

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

Rebound effects of livestock production expansion.

egg: 29 Tg) and 53 Tg of ruminant livestock products (milk: 42 Tg; beef: 6 Tg; lamb: 4 Tg) in 2014. We estimated that 226 Tg food waste (equivalent to 54 Tg in dry matter; 7 Tg in crude protein; 690 billion MJ in energy) and 163 Tg food processing by-products (equivalent to 139 Tg in dry matter; 49 Tg in crude protein; 1907 billion MJ in energy) was available in China in 2014, but only 39% of the food waste and 51% of the food processing by-products were recycled as feed, with the remainder disposed in landfills and incinerators (Supplementary Tables 3-4). The limited use of food waste for feed production in China is primarily due to the early stage of industrialization of recycling food waste as feed, which currently has a low processing capacity 31, and the reliance of industrialized livestock production on concentrate feed 1. In addition, despite being protein-rich, food processing by-products, such as unprocessed oil cakes, contain anti-nutritional factors that may hinder protein absorption by animals. Although fermentation can effectively eliminate these antinutritional factors and enhance digestion and growth performance ³², its limited adoption in China leads to a large amount of these by-products being discarded in landfills or incinerators. Unlike previous studies that considered recycling food waste and food processing by-products as feed to be costless 11-13, we modelled an increasing cost of more recycled food waste and food processing by-products as feed born by monogastric livestock producers and a decreasing cost associated with less food waste and food processing by-products in landfills and incinerators covered by consumers. We found that upcycling 54-100% of food waste and 100% of food processing by-products as feed in scenarios S1 and S2 increased the share of food waste and food processing by-products used as feed within the total feed use by 10-14% in dry matter (Supplementary Fig. 2). Upcycling increased the supply of feed protein by 27-40% and feed energy by 26-39%, and reduced total feed cost (including feed crops, compound feed, food waste, and food processing by-products) for per unit of monogastric livestock production by 2.1-3.0%. This led to a 23-36% increase in monogastric livestock production in S1 and S2 (Fig. 2b). This shift signifies a transition for China from a net importer of monogastric livestock, importing 1% of output in the baseline (S0), to an exporting nation, with 18-25% of output being exported (Fig. 2e). Ruminant livestock production decreased by 3% as the expansion of monogastric livestock reduced the availability of feed crops and compound feed to ruminant livestock (Fig. 2b). To meet domestic demand, ruminant livestock imports rose from 1% of output in the baseline (S0) to 4% (Fig. 2e). Expanded monogastric livestock production raised the demand for primary feed (i.e., feed crops and compound feed), which suprisingly outweighed the reduction in primary feed use by substituting it with food waste and food processing by-products. The overall feed demand for both monogastric and ruminant livestock increased by 17-34% due to a 33-67% rise in feed demand for monogastric livestock (Fig. 3b). The upcycling increased the feed conversion ratio (FCR, the ratio of fresh feed inputs to live weight gain) for monogastric livestock by 0.22-0.62 kg kg⁻¹, but decreased the edible feed conversion ratio (eFCR, the amount of human-edible feedstuffs, i.e., feed crops and compound feed, used for per unit of live weight gain) by 0.11-0.19 kg kg⁻¹, indicating its reduced reliance on human-edible feedstuffs (Supplementary Fig. 3a). Since feed crops and compound feed account for only 12% of ruminant feed (compared to 88% from grass, see Supplementary Fig. 4d), the upcycling had a minor impact on ruminant production and its FCR and eFCR (Supplementary Fig. 3b). The growing demand for crop used as animal feed increased reliance on crop imports, with the import share rising from 11% in the baseline (S0) to 15-19% (Fig. 2d), considering that the total crop production declined by 1.2-4.4% (Fig. 2a). However, the crop cultivated area expanded by 0.6-13% due to the different cultivated area intensities of crops (Fig. 3a). Detailed impacts on crop production structure, as well as the use of N and P fertilisers, were explicitly presented in Supplementary Results. Adjustments in crop and livestock production also had knock-on effects beyond the agricultural

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

sectors in the broader economy, thereby influencing sectoral employment, gross domestic product

(GDP), and household welfare (a measure of economic well-being in US dollars). We observed that the increase of 11.5-18.4 million people in employment in monogastric livestock production was largely a transfer from the non-food sector (i.e., industries and services; detailed in Appendix Table 1) (Supplementary Fig. 7a,c). The non-food sector experienced a slight relative output decline of 1.0-1.4% (Supplementary Fig. 8a,c) and an absolute loss of 28-41 billion US dollars (USD, 2014 constant price) (Supplementary Fig. 9a). In contrast, N and P fertiliser production surged by 35-36% and 20-59% (Fig. 2c), respectively, due to rising demand and decreased production costs, as the shrinking non-food sector made key inputs more available to fertiliser production. As a consequence, China became an exporter of N fertiliser and P fertiliser (Fig. 2f). The absolute value of fertiliser output rose by 5.4-7.0 billion USD (Supplementary Fig. 9a), which compensated less than one-fifth of the total output decrease of the non-food sector. The economic losses in the crop and non-food sectors were largely offset by the expansion of the monogastric livestock and fertiliser sectors (Supplementary Fig. 9a). The overall impact on China's economy was a 0.02-0.07% (0.8-2.6 billion USD) decrease in GDP (Supplementary Fig. 11) and a slight positive impact on household welfare (0.18-0.32%) (Supplementary Fig. 12).

Asymmetric impacts of upcycling food waste and food processing by-products.

We found that the 23-36% expansion in monogastric livestock production in scenarios S1 and S2 increased Chinese economy-wide emissions of acidification polluants by 2.5-4.0% (Fig. 4b), and eutrophication pollutants by $\pm 0.2\%$ (Fig. 4c). The 0.5-1.4% decrease in economy-wide GHG emissions was caused by less food waste and food processing by-products in landfills and incinerators and contraction of the non-food sectors (Fig. 4a). China's main food and feed trading partners (MTP, including Brazil, the United States, and Canada) experienced a reduction in economy-wide emissions of GHGs by 1.1-1.3%, acidification pollutants by 8-13%, and eutrophication pollutants by 2.5-4.0%. These environmental benefits for MTP arose from a reduction in their domestic livestock and fertiliser production, as China shifted from a net importer to an exporter of livestock products and fertilisers (Fig. 2e,f).

For assessing food security, we used four indicators covering two dimensions. Two indicators for food availability, i.e., dietary calorie availability and the population at risk of hunger. Two indicators

for food access, i.e., cereals affordability for labour force and the average food price (including primary food products and processed food). Our findings suggest that upcycling, accompanied by resource reallocation across the whole economy, enhanced food security in China without compromising that of its trading partners. In addition, the reduced cost of collecting food waste and food processing by-products for landfill and incineration enabled consumers in China to allocate more of their income to food consumption. Since the cost of food waste collection for landfill and incineration was quite small in the baseline (S0), the impact of reduced collection costs only had a modest positive effect on most food security indicators. Globally, the average food price declined by 0.1-0.2% (Fig. 5a,e). In China, dietary calorie availability increased by 0.16-0.32%, and the population at risk of hunger, representing 17% of the global population at risk of hunger, decreased by 1.6-3.2% (Fig. 5c,d). Cereals affordability for labour force increased by 0.29-0.47% (Fig. 5b), as a result of a rise in the average wage across the Chinese economy (0.13-0.22%) (Supplementary Fig. 5) and a decrease in cereals price (0.16-0.26%) (Supplementary Fig. 15).

Absorbing rebound effects through emission taxes.

A modest mitigation target of S3 could absorb the rebound effects of upcycling food waste and food processing by-products as feed in China (Fig. 4) and safeguard global food security. Changes in food security indicators under S3 were nearly identical to those in S1 (Fig. 5). This is due to the implementation of a low tax rate on emissions of acidification pollutants (3 \$ ton⁻¹ NH₃-eq) in China. The reduction in emissions of all pollutants in S3 was mainly attributed to a decrease in total crop production compared to S1 (Fig. 2a; Fig 4; Supplementary Fig. 14a,b,c). Livestock production also slightly decreased in scenario S3 (Fig. 2b). However, P fertiliser production increased by 40% while N fertiliser production decreased by 6% compared to S1 (Fig. 2c). As a result, emissions increased in MTP compared to S1 (Fig. 4) due to a shift of emission-intensive production from China to MTP. Nonetheless, emissions of all pollutants in MTP still remained below baseline (S0) levels.

An ambitious emission mitigation target of S4 counteracted the rebound effects further and achieved a further emission reduction, but could pose a risk to food security, as the average global food price increased by 9.4% (Fig. 5a,e) and cereals affordability for labour force decreased by 20% in China (Fig. 5b) and by 15% in MTP (Fig. 5f). The negative impact on food security in China and MTP

was a result of the higher tax rates on emissions in both regions (5 \$ ton⁻¹ CO₂-eq , 788 \$ ton⁻¹ NH₃-eq, and 6969 \$ ton⁻¹ N-eq in China; 2.5 \$ ton⁻¹ CO₂-eq in MTP). Food availability in MTP decreased by 3.3%, while it increased by 3.6% in China (Fig. 5d,h). The latter was a result of consumers transitioning from ruminant-sourced food to less expensive plant and monogastric-sourced food in China (Supplemntary Fig. 16c). Consequently, the population at risk of hunger in MTP increased by 346%, but declined in China by 36% (Fig. 5 c,g). The 2.6% reduction in total GHG emissions and the 2.5% decrease in emissions of acidification pollutants in China in S4 were largely driven by the non-food production contraction compared to S1 (Fig. 4a,b). The 2.0% reduction in total emissions of eutrophication pollutants (Fig. 4c) in China was mainly the result of shifting from ruminant to monogastric livestock production (Supplementary Fig. 14f). For MTP, the 2.0% reduction in total GHG emissions was largely attributed to reductions in total crop and livestock production (Fig. 4a). Meanwhile, emissions of acidification and eutrophication pollutants decreased both by 5% in MTP (Fig. 4b,c).

Discussion

We explored the possible environmental and economic consequences of upcycling food waste and food processing by-products in China's monogastric livestock production in a global context, and provided possible solutions to absorb the rebound effects in China and safeguard global food security. Our study serves as a step towards bridging monetary AGE models with biophysical and nutritional (e.g. protein and energy) constraints. Our integrated environmental-economic framework complements previous linear optimisation studies ¹¹⁻¹³, which overlooked market-mediated effects via the price system. Our modelling framework captured the indirect "rebound effect" of livestock production expansion induced by lower feed costs and its knock-on effects beyond the agricultural sectors, which may undermine the expected environmental benefits in the transition to more circular food systems. Further, we showed that changes in China's food production structure had significant cross-border impacts on its trading partners.

Upcycling food waste and food processing by-products as animal feed.

The primary challenges in upcycling food waste and processing by-products as animal feed are concerns over food and feed safety and potential animal health risks. For example, European Union

(EU) legislation prohibits food waste in animal feed due to disease transmission concerns ³³. In contrast, feeding animals with food waste is more prevalent in Asian countries such as China, South Korea, and Japan, driven by growing demand for animal-sourced food, resource constraints that prioritize food production over feed, and the preference for low-cost alternative feeds among smallscale farms ⁹. Extensive field-based evidence has demonstrated that feeding animals with properly treated food waste is safe for animals with minimal health risks ³⁴. Thermal treatment methods, including heating, drying, and dehydration, are the most commonly used approaches to effectively reduce pathogen transmission risks and ensure food and feed safety 9. While upcycling food waste as feed has been shown not to affect livestock productivity 9, to gain acceptance and adoption among livestock producers, livestock production from food waste must demonstrate its economic competitiveness against conventional feed 34. Upcycling food waste and food processing byproducts as feed necessitates various investments and policies to support the construction of municipal food waste collection plants to efficiently collet, sanitize, and package discarded food waste and food processing by-products for sale to livestock producers as feed ¹². Achieving nearfull use of food waste and food processing by-products as feed appears feasible in China in the future due to several reasons. First, the food waste treatment industry (i.e., food waste collection service and food waste recycling service) has seen significant development and expansion in recent years ³⁵. Second, reinforced policies on municipal solid waste separation and collection guarantee a stable feed supply for monogastric livestock production ³⁶. For example, the Chinese government recently launched an action plan to reduce reliance on soybean imports, which includes a key initiative to give a trial to feed production from food waste in 20 cities by 2025 ³⁷. Additionally, the geographic proximity of industrial livestock farms to municipal food waste collection plants further facilitates the feasibility of upcycling ³⁵.

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

Rebound effects of upcycling food waste and food processing by-products as animal feed.

Policymakers focused on reducing the environmental impact of food systems and enhancing food security may find our findings particularly informative, as we unveil the asymmetric impacts of upcycling food waste and food processing by-products as feed on food security and environment sustainability. On the one hand, rebound effects, where lower feed costs lead to an expansion of monogastric livestock production, diminish the environmental benefits of upcycling food waste and

food processing by-products as feed. We observed Chinese economy-wide emissions of acidification and eutrophication polluants increased by 2.5-4.0% and by ±0.2% in scenarios S1 and S2. In contracst, the 0.5-1.4% decrease in economy-wide GHG emissions was caused by less food waste and food processing by-products in landfills and incinerators and contraction of the non-food sectors. China's trading partners obtained environmental benefits through reducing their domestic livestock and fertiliser production, as China shifted from a net importer to an exporter of livestock products and fertilisers. On the other hand, this upcycling accompanying with resource reallocation across the whole economy enhanced food security in China without compromising that of its trading partners. Our results echo the findings of Hegwood, et al. ¹⁹, who argued that rebound effects could offset more than half of avoided food loss and waste, with reductions in environmental benefits and improvements in food security. Our analysis, thus, enhance the understanding of synergies and trade-offs between economic impacts and multiple environmental stresses associated with upcycling food waste and food processing by-products as feed.

Interconnection between food security and environmental sustainability.

Our study highlights the need to integrate both food security and environmental sustainability into policy decisions to leverage potential win-win opportunities, especially under the current challenges such as climate change and resource constraints. In essence, policymakers should pay closer attention to the interconnection between food security and environmental sustainability to better leverage potential synergies and minimize trade-offs ³⁸. The reduction in GHG emissions, coupled with the enhancements in food security, underscores the rationale for policymakers to promote upcycling food waste and food processing by-products as feed. This also aligns with China's recent emphasis on carbon neutrality and food security as leading priorities ^{39,40}. However, policymakers should remain vigilant regarding indirect effects and spillovers, particularly the unintended increases in emissions of acidification and eutrophication pollutants. We implemented two emission mitigation measures to absorb the rebound effects of upcycling food waste and food processing by-products as feed in China. Our findings revealed that high emission taxes counteracted rebound effects but led to a 9.4% rise in food prices, thereby threatening global food security. This aligns findings of Hasegawa, et al. ²¹, who revealed the risk of increased food insecurity under stringent global climate change mitigation policy. Conversely, modest emission taxes provided an

opportunity to absorb the rebound effects in China and safeguard global food security. Therefore, to avoid unintended negative environmental impacts and achieve the dual dividend of environmental sustainability and food security, it is essential to carefully design and implement tailored, complementary policies and measures rather than relying on a single, one-size-fits-all solution. In China, the responsibility for food security and environmental sustainability falls to different government agencies, highlighting the pressing need for improved coordination and consistency within the government to effectively tackle these intertwined issues 41. In addition, a globally coordinated mitigation policy is imperative for reducing the exceedance of the planetary boundaries, as unilateral environmental policies can lead to 'carbon leakage' by outsourcing the production of emission-intensive goods to countries which lack environmental regulations ⁴². Despite the integrated and holistic approach, our study has some limitations as discussed further in Supplementary Discussion. Our integrated environmental-economic framework supports policy design aimed at achieving the dual dividend of environmental sustainability and food security. Our analysis holds significant policy implications not only for China, a key global market for food and feed, but also serves as a blueprint for other populous emerging economies striving to achieve a better balance between food security and environmental sustainability.

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

Methods

The integrated environmental-economic model and database.

The integrated environmental-economic model based on an AGE framework has been widely used to identify the optimal solution towards greater sustainability and enable efficient allocation of resources in the economy under social welfare maximisation ⁴³⁻⁴⁷. For this study, we developed a global comparative static AGE model, a modified version of an integrated environmental-economic model, ^{42,48-52} and improved the representation of food-related (crop and livestock) sectors and associated non-food (compound feed, food processing by-products, nitrogen and phosphorous fertiliser, food waste treatment, and non-food) sectors (see Fig. 1). While the static model has limitations in short-term policy analysis, it minimises assumptions and uncertainties about future economic conditions about technological and resource changes over time, allowing us to isolate the impact of feeding China's monogastric livestock with food waste and food processing by-products. Our model distinguished two regions: China and its main food and feed trading partners (MTP, including Brazil, the United States, and Canada). These partners accounted for more than 75% of China's total trade volume related to food and feed in 2014. Our reference year is 2014, which represents the latest available year of the Global Trade Analysis Project (GTAP) database. Our model is solved using the general algebraic modelling system (GAMS) software package ⁵³.

Modelling circularity in livestock production requires a detailed representation of biophysical flows to consider nutritional balances and livestock feeding requirement due to increased utilisation of food waste and food processing by-products as feed for monogastric livestock production. Following Gatto, et al. ⁵⁴, we converted dollar-based quantities to physical quantities (Tg) to allow the tracing of biophysical flows through the global economy. GTAP version 10 database ⁵⁵ was used to calibrate our AGE model and provide dollar-based quantities. We designed a sectoral aggregation scheme comprising 16 sectors (see Appendix Table 1) based on the original GTAP database to produce social accounting matrices (SAM) (see Appendix Tables 2-3) in our study. Data on physical quantities (see Supplementary Table 2) of crop and livestock production was obtained from FAO ²⁷. Feed production was extracted from "Feed" in the FAO food balance sheet. Grass from natural grassland was derived from Miao and Zhang ⁵⁶. We only included grass from natural grassland where ruminant livestock is grazing for feed, and grass from remaining grassland was excluded. Data on the trade shares matrix was calculated from the UN Comtrade Database ⁵⁷.

Livestock categories were aggregated into two sectors, i.e., monogastric livestock (including pigs, broilers, and laying hens) and ruminant livestock (including dairy cattle, other cattle, and sheep & goats). Furthermore, the inclusion of animal-specific dietary constraints in our model allowed us to calculate the nutritional balance (crude protein and digestible energy), feed conversion ratios (FCR, the ratio of fresh feed inputs to live weight gain), and edible feed conversion ratio (eFCR, the amount of human-edible feedstuffs, i.e., feed crops and compound feed, used for per unit of live weight gain) ⁵⁸ for each livestock sector. First, we obtained the physical quantities (Tg) of feed protein and energy required to produce the output of livestock. Then, the composition of total feed supplied to each livestock sector is specified. When substituting primary feed (i.e., feed crops and compound feed) in animal diets with food waste and food processing by-products, the total protein and total energy supplies per unit of animal output were kept constant in all scenarios. Our FCRs for ruminant livestock are slightly different from FCRs in the literature, as we did not fully account for hay, crop residues, and roughage-like by-products, but this bias did not affect the impacts of feeding food waste and food processing by-products to monogastric livestock. Further model details, nutritional balance, and detailed composition of animals' diets are available in the Supplementary Information (SI).

Modelling amounts and impacts of food waste and food processing by-products.

In this study, we considered food waste and food processing by-products. Food waste was considered a local resource within China, while food processing by-products could be traded between China and MTP. Food waste refers to discarded food products during distribution and consumption. We only considered plant-sourced food waste because animal-sourced food waste

may pose a risk of pathogen transfer, including foot-and-mouth and classical swine fever ⁵⁹. Food waste was quantified separately for each type of food product using data on food consumption and China-specific food loss and waste fractions ²⁸ following the FAO methodology ⁶⁰. Four types of food waste were distinguished, including cereal grains waste, vegetables & fruits waste, roots & tubers waste, and oilseeds & pulses waste. Food processing by-products refer to by-products produced during the food processing stage, including cereal bran, alcoholic pulp (including distiller's grains from maize ethanol production, brewer's grains from barley beer production, and distiller's grains from liquor production), and oil cakes (including soybean cake and other oil cakes). Food processing by-products were estimated from the consumption of food products and specific technical conversion factors ⁶¹. The total amounts of food waste and food processing by-products and their current use as animal feed and discarded biomass (i.e., landfill and incineration) for China in S0 are presented in Supplementary Table 4.

Our model incorporated two food waste-related sectors, i.e., "food waste collection service" and "food waste recycling service sector recycles food waste and food processing by-products as feed for monogastric livestock production. The food waste collection service sector collects food waste and food processing by-products for landfill and incineration. Waste collection, treatment and disposal activities were included in the 'Waste and water (wtr)' sector in the GTAP database. Food waste generation was added as a margin commodity, similar to how GTAP treated transport costs following Peterson ⁶². Thus, the consumer price of food includes both the market price of food and the cost of collecting food waste and food processing by-products. Consumers spend their income on both consumption of goods and food waste collection services, but they derive utility solely from the consumption of goods. In terms of recycling food waste and food processing by-products as feed, monogastric livestock producer bears the associated cost. By multiplying the quantity of food waste with the unit cost of food waste treatment, we can calculate the value of food waste generation. Physical quantities and prices of food waste recycling service and food waste collection service in China were presented in Supplementary Tables 4-5.

Environmental impact assessment.

Three main environmental impacts of food systems were distinguished, i.e., global warming potential (GWP, caused by greenhouse gas (GHG) emissions, including carbon dioxide(CO₂), methane (CH₄), and nitrous oxide (N₂O) emissions; converted to CO₂ equivalents), acidification potential (AP, caused by pollutants leading to acidification, including ammonia (NH₃), nitrogen oxides (NO_x), and sulphur dioxide (SO₂) emissions; converted to NH₃ equivalents), and eutrophication potential (EP, caused by pollutants leading to eutrophication, including N and P losses; converted to N equivalents). The conversion factors for GWP, AP, and EP were derived from Goedkoop, et al. ⁶³. Data on CO₂, CH₄, and N₂O emissions were obtained from the Climate Analysis Indicators Tool (CAIT) ⁶⁴. All GHG emissions calculations in our model follow the IPCC Tier 2 approach ⁶⁵. We derived NH₃, NO_x, and SO₂ emissions from Liu, et al. ⁶⁶, Huang, et al. ⁶⁷, and Dahiya, et al. ⁶⁸, respectively. We considered NO_x emissions from energy use only, as agriculture's contribution to NO_x emissions is generally small ($\leq 2\%$) ⁶⁹. We used the global eutrophication database of food and non-food provided by Hamilton, et al. ⁷ to obtain data on N and P losses to water bodies.

The total emissions of GHGs, acidification pollutants, and eutrophication pollutants for the food and non-food sectors in the base year were calculated first. Then, we allocated the total emissions to specific sectors according to the shares of emissions per sector in total emissions to unify the emission data from different years. Detailed information about emissions sources across sectors is provided in Appendix Table 4. The sectoral-level emissions as well as the US dollar-based emission intensities of GHGs (t CO₂ equivalents million USD⁻¹), acidification pollutants (t NH₃ equivalents million USD⁻¹), and eutrophication pollutants (t N equivalents million USD⁻¹) are presented in Appendix Tables 5-10. We attributed the environmental impacts between the main (e.g., cereal flour) and joint products (e.g., cereal bran) according to their relative economic values (see Supplementary Table 6).

Two types of land use, i.e., cropland and pastureland, were distinguished. We updated the GTAP data on crop harvested areas using the FAO ²⁷ database. Pastureland was defined as areas where

ruminant grazing occurs. We derived nitrogen and phosphorous fertiliser use by crop types and countries from Ludemann, et al. ⁷⁰.

Food security indicators.

The FAO ⁷¹ defines food security as encompassing four key dimensions: availability (adequate food supply), access (sufficient resources to obtain food), utilisation (nutritious and safe diets), and stability (consistent access to food over time). We focused on the first two dimensions. First, food availability is defined as "calories per capita per day available for consumption". "Population at risk of hunger" refers to the portion of people experiencing dietary energy (calorie) deprivation lasting more than a year following the FAO-based approach ⁷². This approach has been widely used in agricultural economic models to evaluate the risk of food insecurity ^{21,73,74}. In essence, the population at risk of hunger is determined by multiplying the prevalence of undernourishment (PoU) by the total population and is based on dietary energy availability calculated by our model. It is assumed that there is no risk of hunger for high-income countries; consequently, the population at risk of hunger is not applied to the United States and Canada ^{21,73,74}. Second, the access dimension is tied to people's purchasing power, which depends on food prices, dietary habits, and income trends ⁷⁵. We calculated the average food price (including primary food products and processed food), and estimated changes in food affordability by subtracting changes in the average wage across the whole economy from fluctuations in cereal prices.

Definition of scenarios.

To estimate the impacts of increased utilisation of food waste and food processing by-products as animal feed on food security and the environment, we examined five scenarios, including one baseline (S0) scenario representing the economies of China and MTP in 2014, two scenarios involving increased utilisation of food waste and food processing by-products as animal feed, and two scenarios with utilisation of food waste and food processing by-products as animal feed combined with emission mitigation measures. We implemented economy-wide emission taxes under the partial use of food waste and food processing by-products as animal feed (scenario S1), considering the perishability and collection challenges of food waste, as well as the reduced availability of food waste for feed in accordance with SDG 12.3 ("halving food waste") ¹⁴. The latter four scenarios were compared to the 2014 baseline (S0) scenario. The scenarios are further described below and in Supplementary Table 1.

- **S1 Partial use of food waste and food processing by-products as feed.** Scenario S1 investigated the impacts of upcycling partial food waste and food processing by-products as feed (54% of food waste and 100% of food processing by-products for monogastric livestock). Cross-provincial transportation of food waste was not allowed in S1, which limits the maximum utilisation rate of food waste with high moisture content to 54% in China, according to Fang, et al. ¹².
- **S2 Full use of food waste and food processing by-products as feed.** Scenario S2 analysed the impacts of upcycling full food waste and food processing by-products as feed (100% of food waste and 100% of food processing by-products for monogastric livestock). Cross-provincial transportation of food waste was allowed in S2 because we considered that new technology would become available for processing food waste with high moisture content. Economies of scale in food waste recycling were considered in S2; a 1% increase in recycled waste resulted in only a 0.078% rise in recycling costs ⁷⁶. Thus, as production scales up, marginal costs decrease and then stabilise.
 - **S3 S1 + A modest emission mitigation target.** Economy-wide and uniform emission taxes were implemented across all sectors (crop, livestock, and non-food) at the regional level to achieve a modest emission mitigation target, assuming that emissions of GHGs, acidification pollutants, and eutrophication pollutants in both China and MTP do not exceed their baseline (S0) levels. For a given emission mitigation target for each type of pollutant, the AGE model can endogenously determine the emission taxes for various pollutants (expressed in \$ per ton of CO₂ equivalents, \$ per ton of NH₃ equivalents, and \$ per ton of N equivalents). This approach is commonly used in the literature ^{21,22,74,77} and allows to identify the most cost-effective mitigation pathway for achieving a given emission mitigation target.

490 **S4 - S1 + An ambitious emission mitigation target.** Economy-wide and uniform emission taxes were implemented across all sectors (crop, livestock, and non-food) at the regional level to achieve an ambitious emission mitigation target, assuming that emissions of GHGs, acidification pollutants, and eutrophication pollutants remain within the emission thresholds set by China's and the MTP's annual GHG mitigation targets under the Intended Nationally Determined Contributions (INDC) of the Paris Agreement ^{24,25}, as well as China's emission reduction goals for acidification and eutrophication pollutants in line with the "14th Five-Year Plan" ²⁶.

Data availability

497

507

508

509

510

498 The data and parameters that support the economic model in this study are available from the GTAP 499 version 10 database (https://www.gtap.agecon.purdue.edu/databases/v10/). The other data that 500 support splitting food-related (crop and livestock) sectors and associated non-food (compound feed, 501 food processing by-products, nitrogen and phosphorous fertiliser, food waste treatment, and nonfood) sectors from the original database GTAP 10 are publicly available at FAOSTAT 502 UN 503 (http://www.fao.org/faostat/en/#data) and the Comtrade 504 (https://comtrade.un.org/data). The authors declare that all other data supporting the findings of this 505 study are available within the article and its Supplementary Information files, or are available from 506 the corresponding authors upon reasonable request.

Code availability

The authors declare that the GAMS codes for producing the results of this study are available from the corresponding authors upon reasonable request.

References

511

512 1 Bai, Z. et al. China's livestock transition: Driving forces, impacts, and consequences. Science 513 Advances 4, eaar8534 (2018). https://doi.org/doi:10.1126/sciadv.aar8534 514 2 Hu, Y. et al. Food production in China requires intensified measures to be consistent with 515 national and provincial environmental boundaries. Nature Food 1, 572-582 (2020). https://doi.org/10.1038/s43016-020-00143-2 516 517 3 Tilman, D., Balzer, C., Hill, J. & Befort, B. L. Global food demand and the sustainable 518 intensification of agriculture. Proceedings of the national academy of sciences 108, 20260-519 20264 (2011). 520 4 Steinfeld, H. et al. Livestock's long shadow: environmental issues and options. (Food & 521 Agriculture Org., 2006). 522 5 Herrero, M. et al. Greenhouse gas mitigation potentials in the livestock sector. Nature Climate 523 Change 6, 452-461 (2016). https://doi.org/10.1038/Nclimate2925 524 6 Uwizeye, A. et al. Nitrogen emissions along global livestock supply chains. Nature Food 1, 525 437-446 (2020). https://doi.org/10.1038/s43016-020-0113-y 526 7 Hamilton, H. A. et al. Trade and the role of non-food commodities for global eutrophication. 527 *Nature Sustainability* **1**, 314-321 (2018). 528 8 Clark, M. A. et al. Global food system emissions could preclude achieving the 1.5 and 2 C 529 climate change targets. Science 370, 705-708 (2020). 530 9 Wang, Y. et al. Evidence of animal productivity outcomes when fed diets including food waste: 531 A systematic review of global primary data. Resources, Conservation and Recycling 203, 107411 (2024). https://doi.org/https://doi.org/10.1016/j.resconrec.2024.107411 532

533 Gustavsson, J., Cederberg, C., Sonesson, U., Van Otterdijk, R. & Meybeck, A. (FAO Rome, 10 534 2011). 535 11 Van Zanten, H. H. E. et al. Defining a land boundary for sustainable livestock consumption. 536 Global Change Biology **24**, 4185-4194 (2018). https://doi.org/10.1111/gcb.14321 537 12 Fang, Q. et al. Low-opportunity-cost feed can reduce land-use-related environmental impacts 538 by about one-third in China. Nature Food (2023). https://doi.org/10.1038/s43016-023-00813-x 539 13 van Hal, O. et al. Upcycling food leftovers and grass resources through livestock: Impact of 540 livestock system and productivity. Journal of Cleaner Production 219, 485-496 (2019). 541 https://doi.org/https://doi.org/10.1016/j.jclepro.2019.01.329 542 14 Transforming our world: the 2030 agenda for sustainable development, UN. 543 https://sdgs.un.org/2030agenda (2015). 544 15 Ceddia, M. G., Sedlacek, S., Bardsley, N. & Gomez-y-Paloma, S. Sustainable agricultural 545 intensification or Jevons paradox? The role of public governance in tropical South America. 546 Global Environmental Change 23, 1052-1063 (2013). 547 16 Schipper, L. & Grubb, M. On the rebound? Feedback between energy intensities and energy 548 uses in IEA countries. Energy policy 28, 367-388 (2000). 549 17 Sorrell, S., Dimitropoulos, J. & Sommerville, M. Empirical estimates of the direct rebound 550 effect: A review. Energy policy 37, 1356-1371 (2009). 551 18 Gatto, A., Kuiper, M. & van Meijl, H. Economic, social and environmental spillovers decrease 552 the benefits of a global dietary shift. Nature Food (2023). https://doi.org/10.1038/s43016-023-553 00769-y

554 Hegwood, M. et al. Rebound effects could offset more than half of avoided food loss and waste. 19 555 Nature Food 4, 585-595 (2023). https://doi.org/10.1038/s43016-023-00792-z 556 20 Avetisyan, M., Golub, A., Hertel, T., Rose, S. & Henderson, B. Why a Global Carbon Policy 557 Could Have a Dramatic Impact on the Pattern of the Worldwide Livestock Production. Applied 558 Economic **33**, 584-605 (2011).Perspectives and Policy https://doi.org/https://doi.org/10.1093/aepp/ppr026 559 560 21 Hasegawa, T. et al. Risk of increased food insecurity under stringent global climate change 561 mitigation policy. Nature Climate Change 8, 699-703 (2018). https://doi.org/10.1038/s41558-562 018-0230-x 563 22 Jiang, H.-D., Liu, L.-J. & Deng, H.-M. Co-benefit comparison of carbon tax, sulfur tax and 564 nitrogen tax: The case of China. Sustainable Production and Consumption 29, 239-248 (2022). 565 https://doi.org/https://doi.org/10.1016/j.spc.2021.10.017 566 23 Nsabiyeze, A. et al. Tackling climate change in agriculture: A global evaluation of the 567 effectiveness of carbon emission reduction policies. Journal of Cleaner Production 468, 142973 568 (2024). https://doi.org/https://doi.org/10.1016/j.jclepro.2024.142973 569 24 IPCC-WGIII. Summary for policymakers (AR5). (2014). 570 UNFCC. Paris agreement. (2015). 25 571 State Council of the People's Republic of China. Notice of the State Council on Issuing the 26 572 Comprehensive Work Plan for Energy Conservation and Emission Reduction during the 14th Five-Year Plan, https://www.gov.cn/xinwen/2022-01/24/content_5670214.htm (2022). 573 FAO. http://www.fao.org/faostat/en/#data (2022). 574 27

Xue, L. et al. China's food loss and waste embodies increasing environmental impacts. Nature 575 28 576 Food 2, 519-528 (2021). https://doi.org/10.1038/s43016-021-00317-6 577 29 Ministry of Agriculture and Rural Affairs. Guiding Opinions on Promoting Loss Reduction and 578 **Efficiency** Increase in Agricultural Product Processing, 579 http://www.moa.gov.cn/govpublic/XZQYJ/202012/t20201225_6358876.htm (2020). 580 Ministry of Agriculture and Rural Affairs of the People's Republic of China. Technical solution 581 reducing soybean chicken for corn and meal inpig and feed, 582 http://www.moa.gov.cn/gk/nszd 1/2021/202104/t20210421 6366304.htm> (2021). 583 31 Wang, Y. et al. Pursuing zero-grain livestock production in China. One Earth 6, 1748-1758 584 (2023). https://doi.org/https://doi.org/10.1016/j.oneear.2023.10.019 585 32 Mathivanan, R., Selvaraj, P. & Nanjappan, K. Feeding of fermented soybean meal on broiler 586 performance. International Journal of Poultry Science 5, 868-872 (2006). 587 European Commission (EC). No. 1774/2002 of the European Parliament and of the Council of 33 3 October 2002 laying down health rules concerning animal byproducts not intended for human 588 589 consumption. (2002). 590 34 Dou, Z., Toth, J. D. & Westendorf, M. L. Food waste for livestock feeding: Feasibility, safety, 591 implications. Global Food Security and sustainability 17, 154-161 (2018).https://doi.org/https://doi.org/10.1016/j.gfs.2017.12.003 592 593 35 Bai, Z. et al. Investing in mini-livestock production for food security and carbon neutrality in China. Proceedings of the National Academy of Sciences 120, e2304826120 (2023). 594 595 https://doi.org/10.1073/pnas.2304826120

Zhou, M.-H., Shen, S.-L., Xu, Y.-S. & Zhou, A.-N. New policy and implementation of municipal solid waste classification in Shanghai, China. International journal of environmental research and public health 16, 3099 (2019). Ministry of Agriculture and Rural Affairs of China (MARA). Three-year Action Plan for Reducing and Replacing Feed Soybean Meal, https://www.gov.cn/zhengce/zhengceku/2023- 04/14/content 5751409.htm> (2023). Kroll, C., Warchold, A. & Pradhan, P. Sustainable Development Goals (SDGs): Are we successful in turning trade-offs into synergies? Palgrave Communications 5 (2019). Zhang, H. Securing the 'Rice Bowl': China and Global Food Security. (Springer, 2018). Liu, Z. et al. Challenges and opportunities for carbon neutrality in China. Nature Reviews Earth & Environment 3, 141-155 (2022). Lu, Y. et al. Addressing China's grand challenge of achieving food security while ensuring environmental sustainability. Science advances 1, e1400039 (2015). Long, W., Zhu, X., Weikard, H.-P., Oenema, O. & Hou, Y. Exploring sustainable food system transformation options in China: An integrated environmental-economic modelling approach based on the applied general equilibrium framework. Sustainable Production and Consumption , 42-54 (2024). https://doi.org/https://doi.org/10.1016/j.spc.2024.09.004 Keyzer, M. & Van Veen, W. Towards a spatially and socially explicit agricultural policy analysis for China: specification of the Chinagro models. Centre for World Food Studies, Amsterdam, The Netherlands (2005). van Wesenbeeck, L. & herok, C. European and global economic shifts. ENVIRONMENT AND POLICY 45, 138 (2006).

618 Fischer, G. et al. China's agricultural prospects and challenges: Report on scenario simulations 45 619 until 2030 with the Chinagro welfare model covering national, regional and county level. 620 (Centre for World Food Studies, VU University Amsterdam, 2007). 621 Greijdanus, A. Exploring possibilities for reducing greenhouse gas emissions in protein-rich 46 622 food chains MSc. thesis thesis, Wageningen University & Research, (2013). 623 47 Le Thanh, L. Biofuel production in Vietnam: greenhouse gas emissions and socioeconomic 624 impacts Ph.D. thesis thesis, Wageningen University & Research, (2016). 625 48 Zhu, X. & Van Ierland, E. C. Economic Modelling for Water Quantity and Quality Management: 626 A Welfare Program Approach. Water Resources Management 26, 2491-2511 (2012). 627 https://doi.org/10.1007/s11269-012-0029-x 628 Zhu, X., van Wesenbeeck, L. & van Ierland, E. C. Impacts of novel protein foods on sustainable 49 629 food production and consumption: lifestyle change and environmental policy. Environmental 630 and Resource Economics 35, 59-87 (2006). 631 Zhu, X. & Van Ierland, E. C. A model for consumers' preferences for Novel Protein Foods and 50 632 environmental quality. Economic Modelling 22, 720-744 (2005). 633 51 Zhu, X. Environmental-Economic Modelling of Novel Protein Foods: A General Equilibrium 634 Approach, Wageningen University & Research, (2004). 635 Zhu, X. & Van Ierland, E. The enlargement of the European Union: Effects on trade and 52 636 emissions greenhouse **Ecological Economics** 1-14 (2006).of gases. 57, https://doi.org/https://dx.doi.org/10.1016/j.ecolecon.2005.03.030 637 638 53 GAMS. General algebraic modeling system, https://www.gams.com/> (2022).

639 Gatto, A., Kuiper, M., van Middelaar, C. & van Meijl, H. Unveiling the economic and 54 640 environmental impact of policies to promote animal feed for a circular food system. Resources, 641 Conservation Recycling 200, 107317 (2024).and 642 https://doi.org/https://doi.org/10.1016/j.resconrec.2023.107317 643 55 GTAP. GTAP version 10 Database, http://www.gtap.agecon.purdue.edu/ (2014). 644 56 Miao, D. & Zhang, Y. National grassland monitoring report. (2014). 645 UN Comtrade Database. < https://comtrade.un.org/data (2022). 57 646 58 Wilkinson, J. M. Re-defining efficiency of feed use by livestock. *Animal* 5, 1014-1022 (2011). 647 https://doi.org/10.1017/S175173111100005X 648 59 Shurson, G. C. "What a waste"—can we improve sustainability of food animal production 649 systems by recycling food waste streams into animal feed in an era of health, climate, and 650 economic crises? Sustainability 12, 7071 (2020). 651 Gustafsson, J., Cederberg, C., Sonesson, U. & Emanuelsson, A. The methodology of the FAO 60 652 study: Global Food Losses and Food Waste-extent, causes and prevention"-FAO, 2011. (SIK 653 Institutet för livsmedel och bioteknik, 2013). 654 61 FAO. Technical Conversion Factors for Agricultural Commodities. (1997). 655 62 Peterson, E. B. Gtap-m: a gtap model and data base that incorporates domestic margins. GTAP 656 Technical Papers (2006). 657 63 Goedkoop, M. et al. ReCiPe 2008: A life cycle impact assessment method which comprises 658 harmonised category indicators at the midpoint and the endpoint level. 1-126 (2009). Climate Analysis Indicators Tool (CAIT). https://www.climatewatchdata.org/?source=cait 659 64 660 (2014).

661 IPCC. in Agriculture, Forestry and Other Land Use Vol. 4 (Intergovernmental Panel on 65 662 Climate Change, 2006). 663 66 Liu, L. et al. Exploring global changes in agricultural ammonia emissions and their contribution 664 to nitrogen deposition since 1980. Proceedings of the National Academy of Sciences 119, 665 e2121998119 (2022). https://doi.org/doi:10.1073/pnas.2121998119 666 Huang, T. et al. Spatial and Temporal Trends in Global Emissions of Nitrogen Oxides from 667 1960 to 2014. Environmental Science & Technology 51, 7992-8000 (2017). 668 https://doi.org/10.1021/acs.est.7b02235 669 68 Dahiya, S. et al. Ranking the World's Sulfur Dioxide (SO2) Hotspots: 2019–2020. Delhi Center 670 for Research on Energy and Clean Air-Greenpeace India: Chennai, India 48 (2020). 671 Lamsal, L. et al. Application of satellite observations for timely updates to global anthropogenic 69 672 NOx emission inventories. Geophysical Research Letters 38 (2011). 673 70 Ludemann, C. I., Gruere, A., Heffer, P. & Dobermann, A. Global data on fertilizer use by crop 674 and by country. Scientific Data 9, 501 (2022). https://doi.org/10.1038/s41597-022-01592-z 675 71 FAO. Rome Declaration on World Food Security and World Food Summit Plan of Action., 676 (1996). 677 72 FAO. Methodology for the Measurement of Food Deprivation: Updating the Minimum Dietary 678 Energy Requirements. (2008). 679 73 Hasegawa, T., Fujimori, S., Takahashi, K. & Masui, T. Scenarios for the risk of hunger in the 680 twenty-first century using Shared Socioeconomic Pathways. Environmental Research Letters 681 **10**, 014010 (2015).

682	7/4	Hasegawa, T. et al. Consequence of climate mitigation on the risk of hunger. Environmental
683		science & technology 49, 7245-7253 (2015).
684	75	Lele, U. et al. Measuring food and nutrition security: An independent technical assessment and
685		user's guide for existing indicators. Rome: Food Security Information Network, Measuring
686		Food and Nutrition Security Technical Working Group 177 (2016).
687	76	Cialani, C. & Mortazavi, R. The Cost of Urban Waste Management: An Empirical Analysis of
688		Recycling Patterns in Italy. Frontiers in Sustainable Cities 2 (2020).
689		https://doi.org/10.3389/frsc.2020.00008
690	77	Havlík, P. et al. Climate change mitigation through livestock system transitions. Proceedings
691		of the National Academy of Sciences 111, 3709-3714 (2014).
692		
693	Ackno	wledgements
694 695 696 697 698 699 700 701 702	We thank conference participants at III Economy for The Common Good International Conference (ECGIC) and the 29 th Annual Conference of European Association of Environmental and Resource Economists (EAERE) in 2024 for helpful comments and discussions. We acknowledge financial support from the National Natural Science Foundation of China [NSFC, grants no. 32272814], the High-level Team Project of China Agricultural University, and the Agriculture Green Development Program sponsored by China Scholarship Council [no. 201913043]. Artificial Intelligence (in our case ChatGPT) has been used to polish the English writing of paragraphs in this paper. After using this tool/service, we reviewed and edited the content as needed and took full responsibility for the content of the publication.	
703	Autho	r contributions
704 705 706 707	X.Z., H All auth	X.Z., H.P.W., and Y.H. designed the research; W.L. and X.Z. developed the model; W.L., P.W., O.O., and Y.H. analysed data; W.L., X.Z., H.P.W., O.O., and Y.H. wrote the paper nors contributed to the analysis of the results. All authors read and commented on various of the paper.
708	Comp	eting interests
709	The aut	hors declare no competing interests.
710	Additi	onal information

Details about the data, methods, and framework are presented in Supplementary Information (SI).




Fig. 1 | Representation of the economy in China in the applied general equilibrium (AGE) framework with food waste and food processing by-products. The framework includes four parts: (1) Production; (2) Consumption; (3) Food waste generation; (4) Food waste treatment. The generated food waste and food processing by-products are sent either to the 'food waste recycling service' sector or the 'food waste collection service' sector. The food waste recycling service sector recycles food waste and food processing by-products as feed for monogastric livestock production. The food waste collection service sector collects food waste and food processing by-products for landfill and incineration. The consumer price of food includes both the market price of food and the cost of collecting food waste and food processing by-products. The monogastric livestock producer bears the cost of recycling food waste and food processing by-products as feed. Detailed information is presented in Methods and Supplementary Information.




Fig. 2 | Impacts of upcycling food waste and food processing by-products as feed in China's monogastric livestock sector on domestic production and net export of total crop, livestock, and fertiliser. Total (a) crop, (b) livestock, and (c) fertiliser production (Tg) in scenarios. Total (d) crop, (e) livestock, and (f) fertiliser net export (Tg) in scenarios. Total crop production exclude food waste and food processing by-products used by "food waste recycling service" and "food waste collection service" sectors (see Supplementary Table 4 for detailed data). Definitions of scenarios (S1 - 'Partial use of food waste and food processing by-products as feed'; S2 - 'Full use of food waste and food processing by-products as feed'; S3 - 'S1 + A modest emission mitigation target'; S4 - 'S1 + An ambitious emission mitigation target') are described in Table 1.




Fig. 3 | Impacts of upcycling food waste and food processing by-products as feed in China's monogastric livestock sector on domestic total agricultural land use and feed demand. (a) Total agricultural land use (crop harvested area and pastureland) (Mha) and (b) feed demand by monogastric livestock (Tg) in scenarios. Definitions of scenarios (S1 - 'Partial use of food waste and food processing by-products as feed'; S2 - 'Full use of food waste and food processing by-products as feed'; S3 - 'S1 + A modest emission mitigation target'; S4 - 'S1 + An ambitious emission mitigation target') are described in Table 1.




Fig. 4 | Impacts of upcycling food waste and food processing by-products as feed in China's monogastric livestock sector on economy-wide emissions in China (CN) and China's main food and feed trading partners (MTP). Changes in (a) economy-wide emissions of greenhouse gases (Tg CO₂-eq), (b) acidification pollutants (Tg NH₃-eq), and (c) eutrophication pollutants (Tg N-eq) in China and MTP in scenarios with respect to the baseline (S0). MTP includes Brazil, the United States, and Canada. Definitions of scenarios (S1 - 'Partial use of food waste and food processing by-products as feed'; S2 - 'Full use of food waste and food processing by-products as feed'; S3 - 'S1 + A modest emission mitigation target'; S4 - 'S1 + An ambitious emission mitigation target') are described in Table 1.




Fig. 5 | Impacts of upcycling food waste and food processing by-products as feed in monogastric livestock sector on food security indicators in China (CN) and China's main food and feed trading partners (MTP). Changes in (a) average food price (including primary food products and processed food), (b) cereals affordability for labour force, (c) population at risk of hunger (million people; S0 = 140.7 million people), and (d) food availability (kcal capita⁻¹ day⁻¹) in China in scenarios with respect to the baseline (S0). Changes in (e) average food price (including primary food products and processed food), (f) cereals affordability for labour force, (g) population at risk of hunger (million people; S0 = 5.3 million people), and (**d**) food availability (kcal capita⁻¹ day⁻¹) in MTP in scenarios with respect to the baseline (S0). (i) Net imports (Tg) of main food and feed products from MTP to China in the baseline (S0), MTP includes Brazil, the United States, and Canada. According to the FAO approach, it is assumed that there is no risk of hunger for highincome countries; consequently, the population at risk of hunger is not applied to the United States and Canada ^{21,73,74}. Definitions of scenarios (S1 - 'Partial use of food waste and food processing byproducts as feed'; S2 - 'Full use of food waste and food processing by-products as feed'; S3 - 'S1 + A modest emission mitigation target'; S4 - 'S1 + An ambitious emission mitigation target') are described in Table Credit: World 1. Countries base map, Esri (https://hub.arcgis.com/datasets/esri::world-countries/about).

743 744

745

746

747

748

749 750

751

752753

754 755

756

757